arXiv:hep-th/9912098v4 6 Jun 2000

KEK-TH-656
hep-th/9912098
December, 1999

A Candidate for A Renormalizable and Diffeomorphism-Invariant 4D Quantum Theory of Gravity

Ken-ji Hamada[1]

Institute of Particle and Nuclear Studies,
High Energy Accelerator Research Organization (KEK),
Tsukuba, Ibaraki 305-0801, Japan

$\left(\begin{array}{c}\text{revised and extended version}\\ \text{to appear in Prog. Theor. Phys.}\end{array}\right)$

Abstract

We present evidence that there is a 4D model that satisfies the conditions of renormalizability and diffeomorphism invariance simultaneously at the 2-loop level. The traceless mode is treated perturbatively, while the conformal mode can be managed exactly. The two conditions constrain the theory strongly and determine the measure of the gravitational field uniquely. Quantum corrections of the cosmological constant are computed in part to 3-loop diagrams. The method to remove the negative-metric states is also discussed from the viewpoint of diffeomorphism invariance in analogy to the R_ξ gauge in spontaneously broken gauge theory. The model may be a candidate for a continuum version of 4D simplicial quantum geometry realized in recent numerical simulations.

---

[1] E-mail address : hamada@post.kek.jp

1 Introduction

For a long time, many authors have been longing to construct a well-defined 4D quantum theory of gravity [1]–[16]. Renormalizability is a big problem that we must overcome in such a construction. For this reason, we must make a theory of this kind to be diffeomorphism invariant. Furthermore, there are physical problems, such as unitarity, and the correct manner to define the S-matrix in a fully fluctuating geometry. As a first step, it should be clarified whether there is a model that satisfies the two fundamental conditions of renormalizability and diffeomorphism invariance simultaneously. In this paper we give evidence that the model proposed in a previous paper [16] satisfies these two conditions. The problem of the S-matrix and unitarity in our model is also discussed from the viewpoint of diffeomorphism invariance at the end of the paper.

Along another line, there have been developments in the dynamical triangulation approach [17, 18, 19] to a 4-dimensional manifold [20, 21]. Recent numerical simulations suggest that 4D simplicial quantum geometry with $U(1)$ gauge fields is likely to have a phase transition that is of higher order than first [21]. Naively, if there is a continuum limit of 4D simplicial quantum geometry, it should satisfy the conditions of renormalizability and diffeomorphism invariance.

Diffeomorphism invariant quantum theory is formally defined by the functional integration over the metric

$$Z = \int \frac{[g^{-1}dg]_g [df]_g}{\mathrm{vol(diff.)}} \exp[-I(f,g)] \ , \tag{1.1}$$

where f is a matter field [2] and I is an invariant action. The measure of the metric is defined by the norm,

$$< dg, dg >_g = \int d^D x \sqrt{g} g^{\mu\nu} g^{\lambda\sigma} (dg_{\mu\lambda} dg_{\nu\sigma} + u dg_{\mu\nu} dg_{\lambda\sigma}) \ , \tag{1.2}$$

where $u > -1/D$ by the positive definiteness of the norm.

Before the middle 1980s, pertubation theory had been naively defined by replacing the covariant measure with the measure defined on the background metric, $\hat{g}$, as $[g^{-1}dg]_{\hat{g}}[df]_{\hat{g}}$. This replacement, however, breaks the background-metric independence, or, a infinitesimal version of it, diffeomorphism invariance. Renormalizability is also related to the problem of the measure. This problem could not be overcome even by going to supergravity. It has thus been believed that superstring theory [22] is a unique candidate for a background-metric independent quantum theory of gravity [23].

At the end of the 1980s, a new approach emerged. This theory was formulated in such a way as to preserve the background-metric independence, even after the measure is re-expressed in terms of one defined based on the background metric. Two-dimensional quantum gravity was described exactly by Knizhnik, Polyakov and Zamolodchikov (KPZ) [24] and David, Distler and Kawai (DDK) [25] in this way. Subsequently, several 2D models, such as dilaton gravity [26, 27, 28] and R^2-gravity [29], were studied. The idea can apply to even-dimensional quantum gravity, which makes the theory background-metric independent. To do this, we must add an action, S, as in Refs.[16, 26]

---

[2] In this paper all second order fields, except the gravitational field, are referred to as "matter fields".

$$Z = \int \frac{[d\phi]_{\hat{g}}[e^{-h}de^{h}]_{\hat{g}}[df]_{\bar{g}}}{\mathrm{vol(diff.)}} \exp[-S(\phi,\bar{g}) - I(f,g)] \ . \tag{1.3}$$

The metric is now decomposed as $g_{\mu\nu} = e^{2\phi}\bar{g}_{\mu\nu}$ and $\bar{g}_{\mu\nu} = (\hat{g}e^{h})_{\mu\nu}$, where $tr(h) = 0$, adopted by Kawai, Kitazawa and Ninomiya [30] as well as by the present author [16]. The measures of the metric fields are defined on the background metric by the following norms:

$$< d\phi, d\phi >_{\hat{g}} = \int d^D x \sqrt{\hat{g}} (d\phi)^2 \ , \tag{1.4}$$

$$< dh, dh >_{\hat{g}} = \int d^D x \sqrt{\hat{g}} \ tr(e^{-h}de^{h})^2 \ . \tag{1.5}$$

The action, S, is the contribution induced from the measures, and it is related to the conformal anomaly. Note that the existence of an anomaly usually means that a classical symmetry breaks down at the quantum level, but now the anomalous contribution, S, is needed to ensure diffeomorphism invariance at the quantum level.

The action S must satisfy the Wess-Zumino condition [31] to realize background-metric independence, the condition $Z(e^{2\omega}\hat{g}e^{b}) = Z(\hat{g})$, referred to as the BMI condition in the following, where ω and $b^{\mu}{}_{\nu}$ [with $tr(b) = 0$] are local functions. Since the measure $[e^{-h}de^{h}]_{\hat{g}}$, itself, is invariant under the change $e^{h} \to e^{-b}e^{h}$, the BMI condition for the traceless mode represents the condition that $\hat{g}$ and h always appear in the combination $\bar{g} = \hat{g}e^{h}$ in the theory [16]. Although this is a rather trivial condition, it implies that when we calculate S, we can guess the h dependence if the $\hat{g}$ dependence of S can be formed.

In addition to being background-metric independent, namely, invariant under general coordinate transformations,

$$\begin{aligned} \delta\phi &= \frac{1}{D}\hat{\nabla}_{\lambda}\xi^{\lambda} + \xi^{\lambda}\partial_{\lambda}\phi \ , \\ \delta\bar{g}_{\mu\nu} &= \bar{g}_{\mu\lambda}\bar{\nabla}_{\nu}\xi^{\lambda} + \bar{g}_{\nu\lambda}\bar{\nabla}_{\mu}\xi^{\lambda} - \frac{2}{D}\bar{g}_{\mu\nu}\hat{\nabla}_{\lambda}\xi^{\lambda} \ , \end{aligned} \tag{1.6}$$

the theory (1.3) should also be renormalizable. In 4 dimensions, S is a fourth order function. In this paper we assert that the measure of a 4D gravitational field, including S and the fourth order invariant action [5]–[12], are uniquely determined by the two conditions. Thus the integrability condition discussed by Riegert, Fradkin and Tseytlin [32], which is generalized to the form that can be applied to higher loops, should be satisfied. Here, we emphasize that the fourth order terms are required for diffeomorphism invariance, and they are by no means introduced by hand only for renormalizability, as in previous studies [6]–[12].

This paper is organized as follows. In the next section, as an exercise before going to 4D models, we briefly review 2D quantum gravity from the viewpoint of diffeomorphism invariance. The KPZ/DDK formula [24, 25] on a spherical topology is computed using dimensional regularization. In this model, many fundamental items are included. In section 3 we consider the 4D model proposed in a previous paper [16], which is slightly modified here. In 4 dimensions the traceless mode as well as the conformal mode become dynamical. Perturbation theory is applied to the traceless mode, where a dimensionless coupling constant, t, is introduced, while the conformal mode can be managed exactly [30, 16]. The model studied by Antoniadis, Mazur and Mottola [13, 14, 15] as an effective theory of quantum gravity essentially corresponds to our model in the limit $t \to 0$. In this section we present evidence that our model is renormalizable and diffeomorphism invariant, in particular satisfying the integrability conditions discussed in [32] up to order t^2, or the two-loop order for the vacuum polarization of the traceless mode. Higher order renormalizability is also discussed from the viewpoint of diffeomorphism invariance. At the end of this section we discuss our computations of quantum corrections of the cosmological constant up to order t^2, to which diagrams up to 3-loops contribute. The computation is analogous to that for 2D quantum gravity, though the equation used to determine the conformal charge of the cosmological constant becomes biquadratic at this order. It can be solved perturbatively. We finally estimate the strength of the coupling t using the results of recent numerical simulations [21]. Section 4 is devoted to conclusions and discussion. At the end of that section, we discuss proper manner to define the S-matrix and the method to remove the negative-metric states in the fourth order theory required by diffeomorphism invariance. Then, the role of the $\phi\bar{F}$ term in the Wess-Zumino action is emphasized in analogy to the R_ξ gauge in spontaneously broken gauge theory.

Our curvature conventions are $R_{\mu\nu} = R^\lambda{}_{\mu\lambda\nu}$ and $R^\lambda{}_{\mu\sigma\nu} = \partial_\sigma \Gamma^\lambda{}_{\mu\nu} - \cdots$.

2 Review of A Two-dimensional Model

In this section we re-investigate 2D quantum gravity with N massless scalar fields, X [24, 25]. Since we wish to extend these arguments to 4 dimensions, we do not use conformal field theories here. The Wess-Zumino action S in 2 dimensions, referred to as the Liouville action, is given by integrating the 2D conformal anomaly as [34]

$$\begin{aligned} S(\phi,\bar{g}) &= \frac{a}{4\pi}\int d^2x\int_0^\phi d\phi\sqrt{g}R \\ &= \frac{a}{4\pi}\int d^2x\sqrt{\hat{g}}(\bar{g}^{\mu\nu}\partial_\mu\phi\partial_\nu\phi+\bar{R}\phi)\ . \end{aligned} \tag{2.1}$$

Here, the Wess-Zumino condition is defined by

$$S(\phi,\bar{g})=S(\omega,\bar{g})+S(\phi-\omega,e^{2\omega}\bar{g})\ . \tag{2.2}$$

One can easily prove that S satisfies the above equation by dividing the region of integration $[0,\phi]$ into $[0,\omega]$ and $[\omega,\phi]$.

The coefficient a is determined by requiring diffeomorphism invariance. Let us apply the general coordinate transformation (1.6) and $\delta X=\xi^\lambda\partial_\lambda X$ to expression (1.3). The action I is invariant, so that $\delta I(X,g)=0$, but S is not. We obtain the following expression:

$$\delta S(\phi,\bar{g})=\frac{a}{8\pi}\int d^2x\sqrt{\hat{g}}\hat{\nabla}_\lambda\xi^\lambda\bar{R}\ . \tag{2.3}$$

This is easily proved by decomposing the general coordinate transformation into two parts, $\delta=\delta_c+\delta_\omega$, and noting that S is invariant under δ_c, which is a transformation law in which ϕ transforms as if it were a scalar field on the metric $\bar{g}_{\mu\nu}$ such that $\delta_c\bar{g}_{\mu\nu}=\bar{g}_{\mu\lambda}\bar{\nabla}_\nu\xi^\lambda+\bar{g}_{\nu\lambda}\bar{\nabla}_\mu\xi^\lambda$ and $\delta_c\phi=\xi^\lambda\partial_\lambda\phi$, while δ_ω is the conformal change of the metric $\bar{g}_{\mu\nu}$ and the local shift of ϕ defined by $\delta_\omega\bar{g}_{\mu\nu}=2\omega\bar{g}_{\mu\nu}$ and $\delta_\omega\phi=-\omega$, with

$$\omega=-\frac{1}{2}\hat{\nabla}_\lambda\xi^\lambda\ . \tag{2.4}$$

Since ξ is infinitesimal, the r.h.s. of (2.3) can be expressed as $-S(\omega,\bar{g})$ using (2.4). Thus, (2.3) is identically the Wess-Zumino condition (2.2) with an infinitesimal value (2.4).

On the other hand, the measure itself is not invariant, due to the fact that there is no $e^{2\phi}$ factor in the norm, which conformally changes in the form $[d\phi]_{(1+2\omega)\hat{g}}[e^{-h}de^h]_{(1+2\omega)\hat{g}}[dX]_{(1+2\omega)\hat{g}}$, where ω is given by (2.4). This is easily seen by applying the general coordinate transformation (1.6) to the norms defined on the background metric as, for instance,

$$\begin{aligned} &< d\phi', d\phi' >_{\hat{g}}=\int d^2x\sqrt{\hat{g}}(d\phi')^2 \\ &=\int d^2x\sqrt{\hat{g}}(1-\hat{\nabla}_\lambda\xi^\lambda)(d\phi)^2=< d\phi, d\phi >_{(1+2\omega)\hat{g}}\ , \end{aligned} \tag{2.5}$$

where $\phi'=\phi+\delta\phi$, so that $d\phi'=d\phi+\xi^\lambda\partial_\lambda d\phi$. In the second equality of (2.5), the $o(\xi^2)$ term has been omitted.

We first evaluate the measure of the traceless mode. Let us take the gauge-fixing condition $(\hat{P}_1^\dagger \bar{g})_\mu = \hat{\nabla}^\lambda \bar{g}_{\lambda\mu} = 0$ to be as usual, where P_1 is the projection operator to the traceless part such that $\delta\bar{g}_{\mu\nu} = (\bar{P}_1\xi)_{\mu\nu}$. We then obtain

$$\frac{[e^{-h}de^h]_{\hat{g}}}{\mathrm{vol(diff.)}} = [e^{-h}de^h]_{\hat{g}} \det \hat{P}_1^\dagger \bar{P}_1 \delta(\hat{P}_1^\dagger \bar{g}) \ . \tag{2.6}$$

As is wellknown, in two dimensions, the r.h.s. can be further written in the form [34]

$$[d\tau](\det \bar{P}_1^\dagger \bar{P}_1)^{\frac{1}{2}} \ , \tag{2.7}$$

where $[d\tau] = [e^{-h}de^h]_{\hat{g}} \det \hat{P}_1^\dagger \delta(\hat{P}_1^\dagger \bar{g})$ represents integration over the Teichmüller parameter. Note that this rewriting is equivalent to taking the gauge condition as $h^\mu{}_\nu = 0$ [3] up to the zero mode, and hence obtain a gauge-fixing term $iB_{\mu\nu}h^{\mu\nu}$, where $B_{\mu\nu}$ is a symmetric traceless field. The ghost action is then given by $b_{\mu\nu}\delta h^{\mu\nu}|_{\xi^\lambda = c^\lambda}$, which is the well-known bc-ghost system, where $b_{\mu\nu}$ is the anti-ghost related to $B_{\mu\nu}$ and c^λ is the reparametrization ghost. Using the well-known conformal property of the P_1 operator [34], we obtain

$$[e^{-h}de^h]_{(1+2\omega)\hat{g}} = [e^{-h}de^h]_{\hat{g}}\Big(1 + \frac{26}{48\pi}\int d^2x \sqrt{\hat{g}} \hat{\nabla}_\lambda \xi^\lambda \bar{R}\Big) \ . \tag{2.8}$$

The measure of ϕ is easily evaluated by using the bilinear term of the action, $\mathcal{I} = S + I$, as a regulator, which is that of a scalar field. Combined with the results for the traceless mode (2.8) and $N+1$ scalar fields, we obtain

$$\begin{aligned} &[dX]_{(1+2\omega)\hat{g}}[d\phi]_{(1+2\omega)\hat{g}}[e^{-h}de^h]_{(1+2\omega)\hat{g}} \exp(-\mathcal{I}) \\ &\quad = (1 - U(\omega))[dX]_{\hat{g}}[d\phi]_{\hat{g}}[e^{-h}de^h]_{\hat{g}} \exp(-\mathcal{I}) \ , \end{aligned} \tag{2.9}$$

where

$$U(\omega) = -\frac{25-N}{48\pi}\int d^2x \sqrt{\hat{g}} \hat{\nabla}_\lambda \xi^\lambda \bar{R} \ . \tag{2.10}$$

Thus, the variation of the partition function (1.3) w.r.t. the general coordinate transformation becomes

$$\delta Z = \frac{-6a + 25 - N}{48\pi}\Big\langle \int d^2x \sqrt{\hat{g}} \hat{\nabla}_\lambda \xi^\lambda \bar{R} \Big\rangle \ . \tag{2.11}$$

Therefore, the diffeomorphism invariance requires [25]

$$a = \frac{1}{6}(25 - N) \ . \tag{2.12}$$

This is equivalent to the condition that the algebra of the energy-momentum tensor, namely, the Virasoro algebra is closed without central charge at the quantum level.

---

[3] If instead one chooses the ghost determinant as $\det \hat{P}_1^\dagger \bar{P}_1$, one must treat the traceless mode as a quantum field [4, 30].

The lower-dimensional operator generally receives a non-trivial quantum correction. The cosmological constant operator, $\Lambda \int e^{\alpha\phi}$, for instance, where classically $\alpha = \dim[\Lambda] = 2$, recieves the quantum correction

$$\alpha = \dim[\Lambda] = 2 + \gamma_\Lambda \ , \tag{2.13}$$

where γ_Λ is the anomalous dimension. In the following we consider the perturbation theory on α in a flat background as $e^{\alpha\phi} = \sum \frac{\alpha^n}{n!}\phi^n$, and we calculate γ_Λ using dimensional regularization [13, 15].

The combined action in $D = 2 - 2\epsilon$ dimensions is

$$\mathcal{I} = \frac{1}{4\pi}\int d^D x \sqrt{\hat{g}} \Big[a\Big(\bar{g}^{\mu\nu}\partial_\mu\phi\partial_\nu\phi + \bar{R}\phi\Big) + \Lambda e^{\alpha\phi} + \bar{g}^{\mu\nu}\partial_\mu X \partial_\nu X \Big] \ , \tag{2.14}$$

where the bc-ghost is omitted. The conformal mode ϕ and the constant α are now dimensionless, while the field X is of dimension $\mu^{-\epsilon}$, where μ is an arbitrary mass scale. The strategy to compute the anomalous dimension using dimensional regularization is as follows. Consider the coefficient a in front of the Wess-Zumino action to be a bare coupling of dimension $\mu^{-2\epsilon}$. We carry out a renormalization procedure with a being unknown, and compute the anomalous dimension. After regularization we take $\epsilon \to 0$ and substitute the value of a required by diffeomorphism invariance into regularized quantities.

Let us evaluate loop diagrams in the flat background. We can easily see that simple poles come only from the one-loop diagram in Fig. 1. The sum of the two-loop diagram (a) and the associated diagram (b), with the one-loop counterterm insertion shown in Fig. 2, yields only double poles proportional to α^4. The higher-loop diagrams also do not yield simple poles.



Fig. 1: One-loop contribution to the cosmological constant.



(a) (b)

Fig. 2: Two-loop contributions to the cosmological constant. The contribution of a one-loop counterterm is also depicted.

Hence, the renormalization factor defined by the relation $\Lambda = Z_\Lambda \Lambda_r$ is computed as

$$Z_\Lambda = 1 - \frac{\alpha^2}{4\tilde{a}} \frac{1}{\epsilon} + \frac{\alpha^4}{32\tilde{a}^2} \frac{1}{\epsilon^2} + \cdots \ , \tag{2.15}$$

where $\tilde{a}$ is the dimensionless bare coupling defined by $a = \tilde{a}\mu^{-2\epsilon}$. Since the bare coupling, a, should be independent of the arbitrary mass scale, μ, we obtain a finite value in the limit $\epsilon \to 0$:

$$\gamma_\Lambda = \frac{\mu}{Z_\Lambda} \frac{dZ_\Lambda}{d\mu} = \frac{\alpha^2}{2a} \ . \tag{2.16}$$

This value is exact to all orders of the perturbation. We thus obtain the quadratic equation $\alpha = 2 + \frac{\alpha^2}{2a}$. Solving this equation, we obtain

$$\alpha = a - \sqrt{a^2 - 4a} \ , \tag{2.17}$$

where we choose a solution such that $\alpha \to 2$ in the classical limit $a \to \infty$.

We now re-explain diffeomorphism invariance in the context of dimensional regularization. In the curved background we obtain the one-loop exact counterterm

$$\Delta \mathcal{I} = -\frac{25 - N}{48\pi} \frac{\mu^{-2\epsilon}}{\epsilon} \int d^D x \sqrt{\hat{g}} \bar{R} \ . \tag{2.18}$$

From Duff's argument [33], the variation of this counterterm w.r.t. the general coordinate transformation, which is now converted into the conformal change of the background metric, $\delta_\omega \hat{g}_{\mu\nu} = 2\omega \hat{g}_{\mu\nu}$, gives the conformal anomaly $U(\omega)$ (2.10), because the conformal change in the scalar curvature in $D = 2 - 2\epsilon$ dimensions is given by the identity

$$\delta_\omega \sqrt{\hat{g}} \bar{R} = -\epsilon 2\omega \sqrt{\hat{g}} \bar{R} = \epsilon \hat{\nabla}_\lambda \xi^\lambda \sqrt{\hat{g}} \bar{R} \ . \tag{2.19}$$

We thus obtain the value (2.12), as explained above. Substituting this value into expression (2.17), we obtain the well-known KPZ/DDK formula [24, 25] for a spherical topology.

3 Four Dimensional Model

In this section we consider a slightly modified version of a model proposed in Ref. [16]. In 4 dimensions the Wess-Zumino action becomes fourth order, parametrized by the three constants a, b and c in the form

$$\begin{aligned} S(\phi, \bar{g}) &= \frac{1}{(4\pi)^2} \int d^4 x \int_0^\phi d\phi \sqrt{g} \Big[a \Big(F + \frac{2}{3} \Box R \Big) + bG + c \Box R \Big] \\ &= \frac{1}{(4\pi)^2} \int d^4 x \Big[\sqrt{\hat{g}} \Big\{ a\bar{F}\phi + 2b\phi \bar{\Delta}_4 \phi + b \Big(\bar{G} - \frac{2}{3} \bar{\Box} \bar{R} \Big) \phi \Big\} \\ &\qquad - \frac{1}{36} (2a + 2b + 3c) \Big(\sqrt{g} R^2 - \sqrt{\hat{g}} \bar{R}^2 \Big) \Big] \ , \end{aligned} \tag{3.1}$$

where the addition of c is a modification. F is the square of the Weyl tensor, and G is the Euler density. They are defined by

$$F = R_{\mu\nu\lambda\sigma}R^{\mu\nu\lambda\sigma} - 2R_{\mu\nu}R^{\mu\nu} + \frac{1}{3}R^2 \ , \tag{3.2}$$

$$G = R_{\mu\nu\lambda\sigma}R^{\mu\nu\lambda\sigma} - 4R_{\mu\nu}R^{\mu\nu} + R^2 \ . \tag{3.3}$$

The operator Δ_4 is the conformally covariant fourth order operator [32] defined by

$$\Delta_4 = \Box^2 + 2R^{\mu\nu}\nabla_\mu\nabla_\nu - \frac{2}{3}R\Box + \frac{1}{3}(\nabla^\mu R)\nabla_\mu \tag{3.4}$$

and satisfying $\Delta_4 = e^{-4\phi}\bar{\Delta}_4$. Thus, *in order to ensure diffeomorphism invariance in 4 dimensions, the theory must be fourth order in the gravity sector.* Only in that case is the theory renormalizable also.

Here, note that R^2 is not integrable w.r.t. the conformal mode, which is the reason that the number of the independent parameters is three. This fact is related to the integrability condition [32] discussed below.

Let us apply the general coordinate transformation (1.6) to the theory (1.3) in 4 dimensions. The Wess-Zumino action then changes by

$$\delta S(\phi,\bar{g}) = \frac{1}{4(4\pi)^2}\int d^4x\sqrt{\hat{g}}\hat{\nabla}_\lambda\xi^\lambda\Big[a\Big(\bar{F} + \frac{2}{3}\,\bar{\Box}\,\bar{R}\Big) + b\bar{G} + c\,\bar{\Box}\,\bar{R}\Big] \ . \tag{3.5}$$

As in two dimensions, this is simply the Wess-Zumino condition (2.2) for the infinitesimal value

$$\omega = -\frac{1}{4}\hat{\nabla}_\lambda\xi^\lambda \ . \tag{3.6}$$

The measure is also not invariant under the transformation (1.6), which conformally changes in the form $[d\phi]_{(1+2\omega)\hat{g}}[e^{-h}de^h]_{(1+2\omega)\hat{g}}[df]_{(1+2\omega)\bar{g}}$ with (3.6). The coefficients a, b and c are determined by imposing the condition of diffeomorphism invariance, so that the ξ dependences from the action and the measures cancel out.

The equations to determine the coefficients a, b and c are now not so simple. Consider the case that the ξ dependence of the deformed measure can be re-expressed in the form $(1-U(\omega))[d\phi]_{\hat{g}}[e^{-h}de^h]_{\hat{g}}[df]_{\bar{g}}$ using the expression of the conformal anomaly

$$U(\omega) = \frac{1}{(4\pi)^2}\int d^4x\sqrt{\hat{g}}\ \omega\Big[a'\Big(\bar{F} + \frac{2}{3}\,\bar{\Box}\,\bar{R}\Big) + b'\bar{G} + c'\,\bar{\Box}\,\bar{R}\Big] \ . \tag{3.7}$$

The regulator used to evaluate the measures is now the combined action $\mathcal{I} = S + I$, which includes the unknown constants a, b and c, so that the coefficients a', b' and c' become in general functions of a, b and c. Thus, the BMI condition, $U + \delta S = 0$, is now expressed by the equations

$$a'(a,b,c) = a \ , \qquad b'(a,b,c) = b \ , \qquad c'(a,b,c) = c \ . \tag{3.8}$$

To be able to impose this BMI condition it is necessary that the ξ dependence of the measure can indeed be expressed in the above form. This forces the matter fields to be coupled with gravity conformally. As for the gravity sector, we must consider the fourth order invariant action, whose form is restricted by the conditions of renormalizability and diffeomorphism invariance discussed below.

We consider the following invariant action, including fourth order terms: [4]

$$I(A,g) = I_{\rm 4th} + I_{LE} \ , \tag{3.9}$$

where

$$I_{\rm 4th} = \frac{1}{(4\pi)^2}\int d^4x\sqrt{g}\Bigl(\frac{1}{t^2}F + dR^2\Bigr) \ , \tag{3.10}$$

$$I_{LE} = \frac{1}{(4\pi)^2}\int d^4x\sqrt{g}(-m^2R+\Lambda) + I_A(A,g) \ . \tag{3.11}$$

Here, $I_{\rm 4th}$ is the fourth order part, which could be regarded as being a part of the measure. I_{LE} is the usual second order action, which describes low-energy physics. m^2 is the inverse of the gravitational constant, and Λ is the cosmological constant. In this section we consider $U(1)$ gauge fields as an example. I_A is the action of the N_A gauge fields.

Above, we introduced the dimensionless self-coupling constant t only for the traceless mode in the form [30, 16]

$$\bar{g}_{\mu\nu} = (\hat{g}e^{th})_{\mu\nu} = \hat{g}_{\mu\lambda}\Bigl(\delta^\lambda_{\ \nu} + th^\lambda_{\ \nu} + \frac{t^2}{2}(h^2)^\lambda_{\ \nu} + \cdots\Bigr) \ . \tag{3.12}$$

The general coordinate transformation for the traceless mode is then expressed as

$$\begin{aligned} t\delta h^\mu_{\ \nu} &= \hat{\nabla}^\mu\xi_\nu + \hat{\nabla}_\nu\xi^\mu - \frac{1}{2}\delta^\mu_{\ \nu}\hat{\nabla}_\lambda\xi^\lambda + t\xi^\lambda\hat{\nabla}_\lambda h^\mu_{\ \nu} \\ &\quad + \frac{t}{2}h^\mu_{\ \lambda}\Bigl(\hat{\nabla}_\nu\xi^\lambda - \hat{\nabla}^\lambda\xi_\nu\Bigr) + \frac{t}{2}h^\lambda_{\ \nu}\Bigl(\hat{\nabla}^\mu\xi_\lambda - \hat{\nabla}_\lambda\xi^\mu\Bigr) + \cdots \ , \end{aligned} \tag{3.13}$$

where $\xi_\mu = \hat{g}_{\mu\lambda}\xi^\lambda$. The BMI condition (3.8) can then be solved using a perturbation in t.

In the following we consider a model with [16]

$$d = \frac{1}{36}(2a+2b+3c) \ . \tag{3.14}$$

This condition implies that the self-interaction terms of the conformal mode, namely R^2 terms in the combined action, cancel out. This model is analogous to the 2D quantum gravity discussed in section 2. In this paper we assert that *only in this case does the theory become renormalizable and diffeomorphism invariant simultaneously in the perturbation of the traceless mode, where the coefficients a, b and c are uniquely determined by the diffeomorphism invariance.*

---

[4] Note that the coefficient d corresponds to c in Ref.[16].

3.1 Renormalization

To confirm the above assertion, we investigate the renormalizability and diffeomorphism invariance of our model focusing on how crucial condition (3.14) is. We use dimensional regularization to regularize our model as a theory on the metric $\bar{g}_{\mu\nu}$. Here, we assume the renormalizability of our model and derive general formulae for renormalization and diffeomorphism invariance. In the following subsections, we show that no inconsistency appears in our model, at least up to order t^2.

In this paper we consider the $m^2 = 0$ case for simplicity. Thus, the bare combined action becomes

$$\mathcal{I} = \frac{1}{(4\pi)^2}\int d^D x\sqrt{\hat{g}}\Big[2b\phi\bar{\Delta}_4\phi + a\bar{F}\phi + b\Big(\bar{G} - \frac{2}{3}\,\bar{\Box}\,\bar{R}\Big)\phi + \frac{1}{t^2}\bar{F} + \frac{1}{36}(2a+2b+3c)\bar{R}^2 + \Lambda e^{\alpha\phi}\Big] + I_A(A,\bar{g}) \ , \quad (3.15)$$

where $D = 4 - 2\epsilon$. The constants a, b and c are considered to be bare couplings of dimension $\mu^{-2\epsilon}$, and t is the bare coupling of dimension μ^{ϵ}. For later use we introduce the dimensionless bare quantities $\tilde{a}$, $\tilde{b}$, $\tilde{c}$ and $\tilde{t}$, defined by the relations $a = \tilde{a}\mu^{-2\epsilon}$, $b = \tilde{b}\mu^{-2\epsilon}$, $c = \tilde{c}\mu^{-2\epsilon}$ and $t = \tilde{t}\mu^{\epsilon}$. Here, we set the dimension of the conformal mode to zero, while the traceless mode is of dimension $\mu^{-\epsilon}$.

The renormalizations of the quantum fields and couplings are defined by the following relations:

$$\phi = Z_\phi^{1/2}\phi_r \ , \qquad h^\mu{}_\nu = Z_h^{1/2}h^\mu_{r\nu} \quad (3.16)$$

and

$$a = Z_a a_r \ , \qquad b = Z_b b_r \ , \qquad c = Z_c c_r \ , \qquad t = Z_t t_r \ . \quad (3.17)$$

Since the quantum traceless mode must be gauge-fixed to carry out calculations, we introduce here a more well-behaved variable, the background field [35] for the traceless mode defined by

$$\hat{g}_{\mu\nu} = (e^{t\hat{h}})_{\mu\nu} \ , \quad (3.18)$$

where $\hat{h}^\mu{}_\mu = 0$. The renormalization of the background traceless mode is also defined by the relation [35]

$$\hat{h}^\mu{}_\nu = Z_{\hat{h}}^{1/2}\hat{h}_r{}^\mu{}_\nu \ . \quad (3.19)$$

On the other hand, since the dynamics of gravitational fields are governed by fourth order derivative theory, and matter fields are coupled to gravitational fields conformally, the counterterms for the background field become fourth order:

$$\Delta\mathcal{I}_F = -P_F\frac{\mu^{-2\epsilon}}{(4\pi)^2}\int d^D x\sqrt{\hat{g}_r}\hat{F}_r \ , \quad (3.20)$$

$$\Delta\mathcal{I}_G = -P_G\frac{\mu^{-2\epsilon}}{(4\pi)^2}\int d^D x\sqrt{\hat{g}_r}\hat{G}_r \ , \quad (3.21)$$

$$\Delta\mathcal{I}_{R^2} = -P_{R^2}\frac{\mu^{-2\epsilon}}{(4\pi)^2}\int d^D x\sqrt{\hat{g}_r}\hat{R}_r^2 \ . \quad (3.22)$$

Here $\hat{F}_r$, $\hat{G}_r$ and $\hat{R}_r^2$ are the square of the Weyl tensor, the Euler density, and the square of the scalar curvature written in terms of renormalized quantities, respectively. The pole terms, P_F, P_G and P_{R^2}, are expanded as

$$P_F = \frac{f_0}{\epsilon} + \tilde{t}_r^2 \left(\frac{f_1}{\epsilon^2} + \frac{f_1'}{\epsilon} \right) + \cdots \ , \tag{3.23}$$

$$P_G = \frac{g_0}{\epsilon} + \tilde{t}_r^2 \left(\frac{g_1}{\epsilon^2} + \frac{g_1'}{\epsilon} \right) + \cdots \ , \tag{3.24}$$

$$P_{R^2} = \frac{l_0}{\epsilon} + \tilde{t}_r^2 \left(\frac{l_1}{\epsilon^2} + \frac{l_1'}{\epsilon} \right) + \cdots \ , \tag{3.25}$$

where $\tilde{t}_r$ is the dimensionless renormalized coupling. The coefficients f_n, g_n, l_n, and so on, are generally functions of the dimensionless renormalized couplings, $\tilde{a}_r$, $\tilde{b}_r$ and $\tilde{c}_r$.

The counterterms, $\Delta\mathcal{I}_F$ and $\Delta\mathcal{I}_G$, are conformally invariant at $D = 4$. In particular, $\Delta\mathcal{I}_G$ becomes topological, which is just an analog of the counterterm (2.18) in 2 dimensions. In $D = 4 - 2\epsilon$ dimensions, the square of the Weyl tensor and the Euler density satisfy the following identities [33]:

$$\delta_\omega \sqrt{\hat{g}_r} \hat{F}_r = -2\epsilon\omega \sqrt{\hat{g}_r} \left(\hat{F}_r + \frac{2}{3} \Box \hat{R}_r \right) \ , \tag{3.26}$$

$$\delta_\omega \sqrt{\hat{g}_r} \hat{G}_r = -2\epsilon\omega \sqrt{\hat{g}_r} \hat{G}_r \ . \tag{3.27}$$

On the other hand, the counterterm $\Delta\mathcal{I}_{R^2}$ is not conformally invariant at $D = 4$, because $\hat{R}^2$ satisfies the identity

$$\delta_\omega \sqrt{\hat{g}_r} \hat{R}^2 = -12\omega \sqrt{\hat{g}_r} \Box \hat{R} \ . \tag{3.28}$$

This counterterm is related to the renormalizations of the couplings a, b and c, and also the integrability condition of the Wess-Zumino action discussed in [32], which is discussed in detail in section 3.3.

The renormalization of the background traceless mode is now defined by the relation

$$\frac{1}{\tilde{t}^2} \hat{F} = \frac{1}{\tilde{t}_r^2} \hat{F}_r - P_F \hat{F}_r \ . \tag{3.29}$$

We thus obtain the renormalization factor of the coupling,

$$(Z_t)^{-2} = 1 - P_F \tilde{t}_r^2 \ , \tag{3.30}$$

and the relation

$$Z_t Z_{\hat{h}}^{1/2} = 1 \ . \tag{3.31}$$

Therefore, the background metric is not renormalized, and hence $\hat{g}_{r\mu\nu} = \hat{g}_{\mu\nu}$.

3.2 Some results on renormalization factors

We present here some results on the renormalization factors discussed in the previous subsection. Let us first compute the coefficients f_0, g_0 and l_0 in the counterterms. They have already been calculated in Refs.[14, 16] by considering the case $t \to 0$ in a curved background.[5] The interaction terms among quantum fields then vanish, and the kinetic terms reduce to conformal invariant ones due to condition (3.14), so that

$$l_0 = 0 \ . \tag{3.32}$$

The coefficients f_0 and g_0 are given by the values

$$f_0 = -\frac{N_A}{20} - \frac{197}{60} \ , \qquad g_0 = \frac{31 N_A}{360} + \frac{769}{360} \ . \tag{3.33}$$

From this we obtain $Z_t = 1 + \frac{f_0 \tilde{t}_r^2}{2\epsilon} + O(\tilde{t}_r^4)$, and hence the β-function

$$\beta = -\epsilon \tilde{t}_r - \tilde{t}_r \frac{\mu}{Z_t} \frac{dZ_t}{d\mu} = f_0 t_r^3 + \cdots \ . \tag{3.34}$$

Since f_0 is negative, this model is asymptotically free [10, 9].

Next, we see that the Wess-Zumino action is likely to be unrenormalized to order t_r^2. First, consider the self-energy diagrams of the conformal mode in a flat background. The combined action is expanded as

$$\mathcal{I} = \frac{1}{(4\pi)^2} \int d^D x \Big(\mathcal{L}_{\rm kin} + \mathcal{L}_{\rm int}^{(2)} + \mathcal{L}_{\rm int}^{(3)} + \mathcal{L}_{\rm int}^{(4)} + \cdots \Big) \ . \tag{3.35}$$

The kinetic terms for the quantum gravitational fields are [6]

$$\mathcal{L}_{\rm kin} = 2b \ \phi \Box^2 \phi + \frac{1}{2} h_{\mu\nu} \Box^2 h^{\mu\nu} - \partial^\mu \chi^\nu \partial_\mu \chi_\nu + \Big(\frac{1}{3} + dt^2 \Big) \Big(\partial_\lambda \chi^\lambda \Big)^2 \ , \tag{3.36}$$

where $\chi^\mu = \partial^\lambda h^\mu{}_\lambda$, and d is given by (3.14). In this subsection, we take $\Box = \partial^\lambda \partial_\lambda$ for simplicity. The interaction terms used here are

$$\mathcal{L}_{\rm int}^{(2)} = -\frac{2}{3} bt \Box \phi \partial_\mu \partial_\nu h^{\mu\nu} \ , \tag{3.37}$$

$$\begin{aligned} \mathcal{L}_{\rm int}^{(3)} = 2bt \Big(& 2\partial_\mu \phi \partial_\nu \Box \phi + \frac{4}{3} \partial_\mu \partial_\lambda \phi \partial_\nu \partial^\lambda \phi \\ & -\frac{2}{3} \partial^\lambda \phi \partial_\mu \partial_\nu \partial_\lambda \phi - 2 \partial_\mu \partial_\nu \phi \Box \phi \Big) \ h^{\mu\nu} \ , \end{aligned} \tag{3.38}$$

$$\begin{aligned} \mathcal{L}_{\rm int}^{(4)} = 2bt^2 \Big(& \Box \phi \partial_\mu \partial_\nu \phi h^\mu{}_\lambda h^{\lambda\nu} + \partial_\mu \partial_\nu \phi \partial_\lambda \partial_\sigma \phi h^{\mu\nu} h^{\lambda\sigma} \\ & + \text{terms including } \partial h \ \Big) \ , \end{aligned} \tag{3.39}$$

---

[5] In this calculation we do not specify the background metric to be in the form (3.18).

[6] Here, we use the relation $\bar{F} = \bar{G} + 2(\bar{R}^{\mu\nu} \bar{R}_{\mu\nu} - \frac{1}{3} \bar{R}^2)$. The second combination of r.h.s. gives the kinetic term of the traceless mode.

where $\mathcal{L}_{\rm int}^{(2)}$ is derived from the term $-\frac{2}{3}b \,\bar{\Box}\, \bar{R}\phi$. The vertices $\mathcal{L}_{\rm int}^{(3)}$ and $\mathcal{L}_{\rm int}^{(4)}$ are from $2b\phi\bar{\Delta}_4\phi$. These vertices are, of course, decomposed into the renormalized ones and their counterterms.

The propagator of the conformal mode, which is described by a line, is given by

$$\frac{(4\pi)^2}{4\tilde{b}_r}\frac{\mu^{2\epsilon}}{k^4} \ . \tag{3.40}$$

For the traceless mode we consider the gauge-fixing term

$$\mathcal{L}_{GF} = -\frac{1}{\zeta}\chi^\mu N_{\mu\nu}\chi^\nu \ , \tag{3.41}$$

where

$$N_{\mu\nu} = \Box\delta_{\mu\nu} - \left(\frac{1}{3} + dt^2\right)\partial_\mu\partial_\nu \ . \tag{3.42}$$

Here, we take the Feynman-type gauge $\zeta = 1$ in which the gauge-fixing term cancels out the last two of the kinetic terms in (3.36). Then, the propagator of the traceless mode, which is described by the wavy line, becomes

$$(4\pi)^2\frac{1}{k^4}(I_H)^{\mu\nu}{}_{,\lambda\sigma} \ , \tag{3.43}$$

where I_H is the projection operator to the traceless mode, defined by

$$(I_H)^{\mu\nu}{}_{,\lambda\sigma} = \delta^\mu_{(\lambda}\delta^\nu_{\sigma)} - \frac{1}{D}\delta^{\mu\nu}\delta_{\lambda\sigma} \ , \tag{3.44}$$

which satisfies $I_H^2 = I_H$. The ghost action is given by applying the general coordinate transformation to the gauge-fixing condition, which is not coupled with the conformal mode.

Diagrams (a) and (b) in Fig. 3 make contributions of order t_r^2 to $Z_b Z_\phi$, while the diagram in Fig. 4, for example, does not contribute, because such a diagram now becomes order t_r^4, where the vertices in Fig. 4 come from the terms linear in ϕ. In general, diagrams which make contributions proportional to $\tilde{a}_r^k\tilde{b}_r^l\tilde{c}_r^m\tilde{t}_r^{2n}$ to renormalization factors have $n-k-l-m$ loops.



Fig. 3: One-loop contributions of order t_r^2 to Z_ϕ.



Fig. 4: One-loop, but the t_r^4-order contribution to Z_ϕ.

The IR divergence is regularized by introducing an infinitesimal mass, λ, for the traceless mode as $1/k^4 \to 1/(k^2+\lambda^2)^2$. After taking contractions of the tensor indices, we obtain the following contribution from the diagram (a) in Fig. 3:

$$\begin{aligned}
&\int \frac{d^Dk}{(2\pi)^D}\phi_r(k)\phi_r(-k)\Big[-\frac{1}{6}b_r t_r^2 \int \frac{d^Dl}{(2\pi)^D}\frac{1}{l^4\ ((l+k)^2+\lambda^2)^2} \\
&\quad\times\Big\{6(l^2k^6+l^6k^2)+24l^4k^4-16(l\cdot k)(l^2k^4+l^4k^2)-20(l\cdot k)^2l^2k^2 \\
&\qquad -2(l\cdot k)^2(l^4+k^4)+8(l\cdot k)^3(l^2+k^2)+8(l\cdot k)^4 \\
&\qquad +\frac{4-D}{3D}\Big(-36l^4k^4+24(l\cdot k)(l^2k^4+l^4k^2)+40(l\cdot k)^2l^2k^2 \\
&\qquad\qquad -4(l\cdot k)^2(l^4+k^4)-16(l\cdot k)^3(l^2+k^2)-16(l\cdot k)^4\Big)\Big\}\Big] \ . \qquad (3.45)
\end{aligned}$$

Integrating over l in the brackets, we obtain the divergence in the form

$$\frac{2b_r}{(4\pi)^2}\ k^4\Big[-3\tilde{t}_r^2\Big(\frac{1}{\bar\epsilon}-\log\frac{\lambda^2}{\mu^2}+\frac{1}{6}\Big)\Big] \ , \qquad (3.46)$$

where $\frac{1}{\bar\epsilon}=\frac{1}{\epsilon}-\gamma+\log 4\pi$. Here, we also compute finite terms for two-loop computations given below. The non-local term, $\log k^2/\mu^2$, does not appear in the finite term.

Diagram (b) in Fig. 3 is rather simple. The two terms in the vertex $\mathcal{L}_{\rm int}^{(4)}$, shown in (3.39), produce logarithmic divergences. The other terms, including derivatives of h, give vanishing contributions. Hence, we obtain the following divergence:

$$\frac{2b_r}{(4\pi)^2}\ k^4\Big[3\tilde{t}_r^2\Big(\frac{1}{\bar\epsilon}-\log\frac{\lambda^2}{\mu^2}-\frac{5}{12}\Big)\Big] \ . \qquad (3.47)$$

Therefore, the UV divergences as well as the IR divergences cancel out in the sum of (3.46) and (3.47), and we obtain

$$Z_bZ_\phi = 1 \qquad (3.48)$$

at order t_r^2.

This result is related to the counterterm, $\Delta\mathcal{I}_{R^2}$. Since $\sqrt{\hat g}\hat R^2$ produces the kinetic term $\sigma\Box^2\sigma$ when the background field of the conformal mode $\hat g_{\mu\nu}=e^{2\sigma}\delta_{\mu\nu}$ is considered, background-metric independence for the conformal mode implies the relations $Z_aZ_\sigma=Z_bZ_\sigma=Z_cZ_\sigma$ of the renormalization factors, where Z_σ is the renormalization factor of the background conformal mode. Since the background metric is not renormalized such that $Z_\sigma=1$, we obtain

$$Z_\phi = 1 \ , \qquad Z_a=Z_b=Z_c \ . \qquad (3.49)$$

Thus, result (3.48) leads to

$$Z_a=Z_b=Z_c=1 \qquad (3.50)$$

at order t_r^2. Since we do not introduce a coupling for the conformal mode, this result implies that the terms of order t_r^2 in P_{R^2}, (3.25), vanish, so that

$$l_1 = l_1' = 0 \ . \tag{3.51}$$

Here, $l_1 = 0$ is rather trivial, because only the one-loop diagrams contribute to order t_r^2, which is a consequence of the cubic and quartic self-interactions of ϕ being dropped, by condition (3.14). The result (3.51) is evidence that an action S satisfying the integrability condition [32] exists at $t_r \neq 0$. This is discussed in further detail in the next subsection.

The relation (3.31) to order t_r^2 implies that the sum of the vertex diagrams (a) through (e), shown in Fig. 5, makes a finite contribution. This is likely to be true, as in the case of the self-energy diagrams. However, because direct calculations would be quite tedious, we need software to take contractions of the tensor indices into consideration.



Fig. 5: One-loop vertex diagrams to order t_r^2.

Quantum corrections proportional to ϕ^3 and ϕ^4, whose terms are absent in the action $\mathcal{I}$, arise at order t_r^4. Replacing the wavy external lines of diagrams (a) and (b) in Fig. 5 with lines, we obtain corrections to ϕ^3 of order t_r^4. Their sum is also likely to be finite, as in the self-energy diagrams of ϕ. In the same way, the divergences proportional to ϕ^4 may cancel out.

Hence, we see that, from (3.31) and (3.48), the vertices in the Wess-Zumino action are likely to be unrenormalized to order t_r^2, which implies that non-local terms related to such vertices do not appear, and thus the Wess-Zumino action does not change form.

3.3 Diffeomorphism invariance at two-loop

In this subsection we discuss the condition of diffeomorphism invariance at the two-loop level. To clarify the origin of the interactions, we divide the combined Lagrangian into three parts, as follows:

$$\mathcal{L}^C = 2b\phi\bar{\Delta}_4\phi + \frac{1}{t^2}\bar{F} + \mathcal{L}_A(A, \bar{g}) \ , \tag{3.52}$$

$$\mathcal{L}^{CS} = a\bar{F}\phi + b\bar{G}\phi \ , \tag{3.53}$$

$$\mathcal{L}^{NC} = -\frac{2}{3}b \,\bar{\Box}\, \bar{R}\phi + \frac{1}{36}(2a + 2b + 3c)\bar{R}^2 \ . \tag{3.54}$$

Here, $\mathcal{L}^C$ is conformally invariant. Expanding $\mathcal{L}^{CS}$ produces conformally invariant vertices linear in ϕ. On the other hand, the last one, $\mathcal{L}^{NC}$, yields non-conformally invariant vertices.

The regularized 1PI effective action is divided into three parts as

$$\Gamma_{reg}(\phi_r, \bar{g}_r) = \mathcal{I}_r(\phi_r, \bar{g}_r, A_r) + V_{reg}(\phi_r, \bar{g}_r) + W_{reg}(\bar{g}_r, A_r) \ . \tag{3.55}$$

Here, finite corrections V_{reg} and W_{reg} come from loop diagrams. The former represents corrections to the Wess-Zumino action, and the latter represents corrections independent of ϕ, which are related to the counterterms for the traceless mode, $\Delta\mathcal{I}_{F,\ G,\ R^2}$, and those for matter fields. From the sum of the results (3.46) and (3.47), we obtain $V_{reg} = -\frac{7}{4}t_r^2 2b_r\phi\bar{\Delta}_4\phi + \cdots$. As discussed in subsection 3.2, there is no non-local correction in this part.

Diffeomorphism invariance now implies that the 1PI regularized effective action satisfies the equation $\delta\Gamma_{reg} = 0$, or

$$\begin{aligned} \delta_\omega W_{reg}(\bar{g}_r, A_r) \ &= \ \frac{1}{(4\pi)^2}\int d^4x\sqrt{\hat{g}_r}\omega\Big[a_r(1+v_a)\Big(\bar{F}_r + \frac{2}{3}\,\bar{\Box}_r\,\bar{R}_r\Big) \\ &\qquad\qquad + b_r(1+v_b)\bar{G}_r + c_r\,\bar{\Box}_r\,\bar{R}_r\Big] \ , \end{aligned} \tag{3.56}$$

where ω is an infinitesimal value (3.6). The r.h.s. of (3.56) comes from the variation of the actions $\mathcal{I}_r$ and V_{reg}, where v_a and v_b denote contributions from δV_{reg}. While v_b has been computed to order t_r^2, with

$$v_b = -\frac{7}{4}t_r^2 \ , \tag{3.57}$$

v_a has not. The l.h.s. of (3.56) corresponds to the anomalous contributions from the measures $U(\omega)$. Here, $\delta W_{reg} = \delta_\omega W_{reg}$, because W_{reg} is not include the conformal mode ϕ, so that decomposing $\delta\bar{g}_{\mu\nu}$ into $\delta_c\bar{g}_{\mu\nu} = \bar{g}_{\mu\lambda}\bar{\nabla}_\nu\xi^\lambda + \bar{g}_{\nu\lambda}\bar{\nabla}_\mu\xi^\lambda$ and $\delta_\omega\bar{g}_{\mu\nu} = 2\omega\bar{g}_{\mu\nu}$, W_{reg} is invariant under $\delta_c\bar{g}_{\mu\nu}$. As seen below, (3.56) uniquely determines coefficients a_r, b_r and c_r.

The background-metric independence of the traceless mode implies that $\hat{g}$ and h always appear in the combination $\bar{g} = \hat{g}e^h$. Therefore, it is enough to know the $\hat{g}$ dependence of W_{reg} to solve the condition of diffeomorphism invariance.

In the following arguments we consider only the $\hat{g}$ dependence of W_{reg}. We here omit tensor indices for simplicity. The Weyl tensor is then described as $\hat{C}$ and $\hat{C}^2 = \hat{F}$, where the subscript r can be dropped, because $\hat{g}_r = \hat{g}$. The Euler density is also written as $\hat{G} = \hat{B}^2$.

Let us consider the W_{reg} part, which is related to three counterterms, $\Delta\mathcal{I}_F$, $\Delta\mathcal{I}_G$ and $\Delta\mathcal{I}_{R^2}$. First, we discuss the one-loop case [33]. At this order, only the conformally invariant Lagrangian, $\mathcal{L}^C$, contributes to the effective action. From a dimensional analysis, the one-loop effective action should have the form [36]

$$\begin{aligned} W^{(1)} &= \frac{\tilde{t}_r^2}{2}\hat{h}_r k^4 \left(\frac{k^2}{\mu^2}\right)^{-\epsilon}\left(\frac{v}{\epsilon} + v'\right)\hat{h}_r + o(\epsilon) \\ &\to v_f\left(\frac{1}{\epsilon}\hat{F} - \hat{C}\log(-\Box/\mu^2)\hat{C}\right) + v'_f\hat{F} \\ &\quad + v_g\left(\frac{1}{\epsilon}\hat{G} - \hat{B}\log(-\Box/\mu^2)\hat{B}\right) + v'_g\hat{G} + o(\epsilon) \ , \end{aligned} \tag{3.58}$$

where v and v' represent v_f and v_g and v'_f and v'_g, related to the conformally invariant counterterms $\Delta\mathcal{I}_F$ and $\Delta\mathcal{I}_G$, respectively. The IR cutoff mass is canceled out. Going to the second line, $(k^2)^{-\epsilon} = 1 - \epsilon\log k^2 + \frac{1}{2!}\epsilon^2\log^2 k^2 + \cdots$ is used. The coefficients v_f and v_g have been computed as $v_f = f_0$ and $v_g = g_0$ in (3.33). Hence, the counterterm can be written as

$$\Delta W^{(1)} = -\frac{f_0}{\epsilon}\hat{F} - \frac{g_0}{\epsilon}\hat{G} \ , \tag{3.59}$$

and the regularized effective action is described as

$$\begin{aligned} W^{(1)}_{reg} &= W^{(1)} + \Delta W^{(1)} \\ &= -f_0\hat{C}\log(-\Box/\mu^2)\hat{C} - g_0\hat{B}\log(-\Box/\mu^2)\hat{B} + v'_f\hat{F} + v'_g\hat{G} \ . \end{aligned} \tag{3.60}$$

The anomalous contribution is given by the variation of regularized effective action with respect to the conformal change as

$$\delta_\omega W^{(1)}_{reg} = a_0 A + b_0 A' \ , \tag{3.61}$$

with

$$a_0 = 2f_0 \ , \qquad b_0 = 2g_0 \ , \tag{3.62}$$

where $A = \omega(\hat{F} + (2/3)\Box\hat{R})$ and $A' = \omega\hat{G}$. To derive this expression we use the following variational formulae [33]:

$$\delta_\omega\hat{F} = -2\epsilon A \ , \qquad \delta_\omega(\hat{C}\log(-\Box)\hat{C}) = -2A + o(\epsilon) \ , \tag{3.63}$$

$$\delta_\omega\hat{G} = -2\epsilon A' \ , \qquad \delta_\omega(\hat{B}\log(-\Box)\hat{B}) = -2A' + o(\epsilon) \ . \tag{3.64}$$

Here, note that using the formulae, we obtain the equation $\delta_\omega W^{(1)} = 0$,[7] so that $\delta_\omega W^{(1)}_{reg} = \delta_\omega \Delta W^{(1)}$, as used in section 2.

Hence, substituting the value (3.61) into the l.h.s. of (3.56), we obtain the values of a_r and b_r at $t_r = 0$ as a_0 and b_0. Here, note that a_0 is always negative and b_0 is positive, and their sum, $d_0 = \frac{1}{18}(a_0 + b_0)$, is positive.

The terms related to the non-conformally invariant counterterm, $\Delta\mathcal{I}_{R^2}$, in the regularized effective action in general have the forms $\hat{R}\log(-\Box/\mu^2)\hat{R}$ and $\hat{R}^2$. The term $\hat{R}\log(-\Box/\mu^2)\hat{R}$ breaks the condition of diffeomorphism invariance, (3.56), and therefore it must vanish. This condition is now equivalent to the vanishing of l_0, (3.32). This implies that in order for the theory to be diffeomorphism invariant, those parts of the action contributing to the one-loop effective action $W^{(1)}$ should be conformally invariant. This is now ensured by condition (3.14) for the gravitational fields. Matter fields must be conformally coupled to gravity; only in this case does the theory become diffeomorphism invariant. Since the variation of the $\hat{R}^2$ term is proportional to $\Box\hat{R}$ (3.28), this term determines the value of c_r. In the one-loop case, however, the conformal invariance is exact, and hence such a non-conformally invariant term vanishes. Thus, the value of c_r at $t_r = 0$ vanishes.

Let us analyze the effective action of order t_r^2 (or order t_r^4 for $Z_{\hat{h}}$). At this order, the non-conformally invariant terms in the action contribute to it. The effective action of order t_r^2 is divided into three parts as

$$W^{(2)} = W^{(2-loop)} + W^{(counter)} + W^{(1-loop)} \ , \tag{3.65}$$

where $W^{(2-loop)}$ represents contributions of two-loop diagrams, which are derived from the conformally invariant Lagrangian $\mathcal{L}^C$. The contribution from associated diagrams with a one-loop counterterm insertion is represented by $W^{(counter)}$. Furthermore, there are contributions of order t_r^2, but they are one-loop diagrams described by $W^{(1-loop)}$, which include the vertices derived from $\mathcal{L}^{CS}$ and $\mathcal{L}^{NC}$.

Since two-loop diagrams have two internal-momentum integrals, from a simple dimensional analysis, we find that $W^{(2-loop)}$ has the following form:

$$\begin{aligned} W^{(2-loop)} &= \frac{\tilde{t}_r^4}{2}\hat{h}_r k^4 \Big[\Big(\frac{k^2}{\mu^2}\Big)^{-2\epsilon} \Big(\frac{x}{\epsilon^2} + \frac{x'}{\epsilon} + x''\Big) \\ &\qquad + \Big(\frac{k^2}{\mu^2}\Big)^{-\epsilon} \Big(\frac{\lambda^2}{\mu^2}\Big)^{-\epsilon} \Big(\frac{y}{\epsilon^2} + \frac{y'}{\epsilon} + y''\Big) \\ &\qquad + \Big(\frac{\lambda^2}{\mu^2}\Big)^{-2\epsilon} \Big(\frac{z}{\epsilon^2} + \frac{z'}{\epsilon} + z''\Big) \Big] \hat{h}_r \ . \end{aligned} \tag{3.66}$$

Here, the dependences of the momentum and the IR cutoff mass, $(k^2)^{-2\epsilon}$, $(k^2\lambda^2)^{-\epsilon}$ and $(\lambda^2)^{-2\epsilon}$, in each term are the features of two-loop diagrams. On the other hand, since the associated counterterm insertion diagrams are one-loop, $W^{(counter)}$ should have the following form:

---

[7] Precisely, we have $\delta_\omega W^{(1)} = o(\epsilon)$, which reflects the fact that the conformal invariance of $\mathcal{L}^C$ is violated by $o(\epsilon)$ in D dimensions.

$$W^{(counter)} = \frac{\tilde{t}_r^4}{2}\hat{h}_r\Big(-\frac{f_0}{\epsilon}\Big)k^4\Big[\Big(\frac{k^2}{\mu^2}\Big)^{-\epsilon}\Big(\frac{d}{\epsilon}+d'\Big)+\Big(\frac{\lambda^2}{\mu^2}\Big)^{-\epsilon}\Big(\frac{e}{\epsilon}+e'\Big)\Big]\hat{h}_r \ . \qquad (3.67)$$

Since the contributions of two-loop diagrams come from the conformally invariant Lagrangian $\mathcal{L}^C$, the related two-loop counterterms becomes conformally invariant. The conditions of renormalizability and IR finiteness give constraints on the coefficients. From the condition that the non-local divergence, $\frac{1}{\epsilon}\log\frac{k^2}{\mu^2}$, cancels in the sum of $W^{(2-loop)}$ and $W^{(counter)}$, we obtain the constraint $-2x-y+f_0d=0$. The cancellation of the IR divergences gives the constraints $y=z=e=0$ and $y'+2z'-f_0e'=0$. Hence, we obtain the following expression:

$$\begin{aligned}
&W^{(2-loop)} + W^{(counter)} \\
&\to \tilde{t}_r^2\Big[-\frac{x_f}{\epsilon^2}\hat{F} - \frac{x_g}{\epsilon^2}\hat{G} + (x'_f - z'_f - f_0d'_f)\frac{1}{\epsilon}\hat{F} + (x'_g - z'_g - f_0d'_g)\frac{1}{\epsilon}\hat{G} \\
&\qquad + x_f\hat{C}\log^2(-\Box/\mu^2)\hat{C} + x_g\hat{B}\log^2(-\Box/\mu^2)\hat{B} \\
&\qquad - (2x'_f + y'_f - f_0d'_f)\hat{C}\log(-\Box/\mu^2)\hat{C} - (2x'_g + y'_g - f_0d'_g)\hat{B}\log(-\Box/\mu^2)\hat{B} \\
&\qquad + (x''_f + y''_f + z''_f)\hat{F} + (x''_g + y''_g + z''_g)\hat{G}\Big] \ . \qquad (3.68)
\end{aligned}$$

The diffeomorphism invariance (3.56) requires $\delta_\omega W^{(2)}_{reg}$ to be proportional to the anomaly forms A and A'. This implies that the finite term including the square of the logarithm should vanish, so that

$$x_f = x_g = 0 \ , \qquad (3.69)$$

which is equivalent to the vanishing of the double pole in the two-loop counterterm. Thus, the renormalization factor of the background traceless mode, $Z_{\hat{h}}$, does not have double poles of order t_r^4. On the other hand, using the relation between the renormalization factors, $Z_tZ_{\hat{h}}^{1/2}=1$ in (3.31), and the finiteness of the β-function, one can easily prove that $Z_{\hat{h}}$ does not have double poles of order t_r^4.[8] Thus, the relation $Z_tZ_{\hat{h}}^{1/2}=1$ guarantees that we can impose the condition of diffeomorphism invariance at the two-loop level.

Furthermore, the relation $Z_t = Z_{\hat{h}}^{-1/2}$ implies that the diagrams with counterterm insertions at vertices and at propagators cancel each other, and hence $W^{(counter)}$ vanishes.[9] The constraint on the IR divergences now becomes $y'_{f,g}+$

---

[8] There is an analogy in QED, where the renormalization factor, defined by $e=Z_ee_r$, satisfies the relation $Z_eZ_3^{1/2}=1$ as a result of the Ward-Takahashi identity $Z_1=Z_2$. The regularized 1PI vacuum polarization does not have $\log^n(-\Box/\mu^2)$ with $n\geq 2$. (See, for instance, section 8-4-4 in Ref.[37].)

[9] Precisely, gauge-fixed renormalization insertion diagrams may be necessary [35].

$2z'_{f,g} = 0$. Therefore, expression (3.68) reduces to the form [10]

$$\tilde{t}_r^2 \Big[(x'_f - z'_f) \frac{1}{\epsilon} \hat{F} + (x'_g - z'_g) \frac{1}{\epsilon} \hat{G} \\ -2(x'_f - z'_f) \hat{C} \log(-\Box/\mu^2) \hat{C} - 2(x'_g - z'_g) \hat{B} \log(-\Box/\mu^2) \hat{B} \\ +(x''_f + y''_f + z''_f) \hat{F} + (x''_g + y''_g + z''_g) \hat{G} \Big] \ . \tag{3.70}$$

Next we consider the one-loop part of the effective action $W^{(1-loop)}$. Recall that it includes the interactions derived from the Lagrangians $\mathcal{L}^{CS}$ and $\mathcal{L}^{NC}$. Since $\mathcal{L}^{NC}$ is not conformally invariant, we must consider here the non-conformally invariant counterterm $\Delta \mathcal{I}_{R^2}$ also. From a dimensional analysis, we obtain the form

$$\begin{aligned} W^{(1-loop)} &= \frac{\tilde{t}_r^4}{2} \hat{h}_r k^4 \Big(\frac{k^2}{\mu^2} \Big)^{-\epsilon} \Big(\frac{w}{\epsilon} + w' \Big) \hat{h}_r \ . \\ &\to \tilde{t}_r^2 \Big[w_f \Big(\frac{1}{\epsilon} \hat{F} - \hat{C} \log(-\Box/\mu^2) \hat{C} \Big) + w'_f \hat{F} \\ &\qquad + w_g \Big(\frac{1}{\epsilon} \hat{G} - \hat{B} \log(-\Box/\mu^2) \hat{B} \Big) + w'_g \hat{G} \\ &\qquad + w_l \Big(\frac{1}{\epsilon} \hat{R}^2 - \hat{R} \log(-\Box/\mu^2) \hat{R} \Big) + w'_l \hat{R}^2 \Big] \ . \end{aligned} \tag{3.71}$$

Combining contributions (3.70) and (3.71), the two-loop counterterm reads

$$\Delta W^{(2)} = -\frac{\tilde{t}_r^2}{\epsilon} \Big(f'_1 \hat{F} + g'_1 \hat{G} + l'_1 \hat{R}^2 \Big) \ , \tag{3.72}$$

with

$$f'_1 = x'_f - z'_f + w_f \ , \qquad g'_1 = x'_g - z'_g + w_g \ , \qquad l'_1 = w_l \ . \tag{3.73}$$

The regularized effective action, $W^{(2)}_{reg} = W^{(2)} + \Delta W^{(2)}$, can then be expressed in the form

$$\begin{aligned} W^{(2)}_{reg} &= t_r^2 \Big[-f''_1 \hat{C} \log(-\Box/\mu^2) \hat{C} - g''_1 \hat{B} \log(-\Box/\mu^2) \hat{B} \\ &\qquad -l'_1 \hat{R} \log(-\Box/\mu^2) \hat{R} + u_f \hat{F} + u_g \hat{G} + u_l \hat{R}^2 \Big] \ , \end{aligned} \tag{3.74}$$

with

$$f''_1 = 2(x'_f - z'_f) + w_f \ , \qquad g''_1 = 2(x'_g - z'_g) + w_g \ , \tag{3.75}$$

where $u_f = x''_f + y''_f + z''_f + w'_f$, $u_g = x''_g + y''_g + z''_g + w'_g$ and $u_l = w'_l$.

The requirement of diffeomorphism invariance is now equivalent to the requirement that the non-local term $\hat{R} \log(-\Box/\mu^2) \hat{R}$ vanishes, or $l'_1 = 0$. This has been indirectly verified by computing the self-energy diagrams of the conformal mode to order t_r^2 in subsection 3.2. From the viewpoint of the vacuum polarization of $\hat{h}_{\mu\nu}$, this implies that the divergences contributing to l_1' in the one-loop diagrams, which include the vertices coming from the non-conformally invariant Lagrangian, $\mathcal{L}^{NC}$, cancel out. As discussed at the end of this subsection, this is consistent with the result in Ref.[16].

---

[10] Note that, from a dimensional analysis, we cannot say that $\delta_\omega W^{(2-loop)}$ vanishes. Rather, it satisfies $\mu \frac{\partial}{\partial \mu} W^{(2-loop)} = 0$.

Furthermore, the non-local term $\phi\Box^2\log(-\Box/\mu^2)\phi$, which changes the form of the Wess-Zumino action, does not appear in the regularized effective action at order t_r^2. The background-metric independence of the conformal mode implies that the vanishing of this term is equivalent to the vanishing of the non-local term $\hat{R}\log(-\Box/\mu^2)\hat{R}$.

The anomalous contribution at the two-loop level is now given by

$$\delta_\omega W_{reg}^{(2)} = t_r^2\Big(a_1 A + b_1 A' + c_1\omega\Box\hat{R}\Big) , \tag{3.76}$$

with

$$a_1 = 2f_1'' , \qquad b_1 = 2g_1'' , \qquad c_1 = -12u_l . \tag{3.77}$$

Combining the one-loop results, (3.56) is now solved as

$$\begin{aligned} a_r(1+v_a) &= a_0 + a_1 t_r^2 + \cdots , \\ b_r(1+v_b) &= b_0 + b_1 t_r^2 + \cdots , \\ c_r &= c_1 t_r^2 + \cdots . \end{aligned} \tag{3.78}$$

The coefficients a_1, b_1 and c_1 are generally functions of the couplings a_r and b_r, while a_0 and b_0 are given by the constants. Hence, the BMI condition can be solved perturbatively. Substituting the value of v_b from (3.57), we then obtain

$$b_r = b_0 + \Big(b_1 + \frac{7}{4}b_0\Big)t_r^2 + \cdots , \tag{3.79}$$

where b_1 is now a function of a_0 and b_0. This equation is used in subsection 3.4.

At higher order, the conditions of diffeomorphism invariance are expressed as the vanishing of the non-local terms: $\hat{C}\log^n(-\Box/\mu^2)\hat{C}$ $(n \geq 2)$, $\hat{B}\log^n(-\Box/\mu^2)\hat{B}$ $(n \geq 2)$ and $\hat{R}\log^n(-\Box/\mu^2)\hat{R}$ $(n \geq 1)$. Each term with $n \geq 2$ is likely to vanish to all orders, as in gauge theories. This is ensured by relation (3.31). The non-trivial condition is the vanishing of the term $\hat{R}\log(-\Box/\mu^2)\hat{R}$, which is the generalized form of the integrability condition discussed by Riegert, Fradkin and Tseytlin [32]. This condition does not imply that $\Delta\mathcal{I}_{R^2}$ vanishes at higher order. This condition may be realized when we substitute the values a_r, b_r and c_r from (3.78) into the expression of the regularized effective action.

This seems to be a trick of dimensional regularization. If there is a regularization defined exactly in 4 dimensions, like the Pauli-Villars method, one would be able to compute the effective action substituting the values of the constants a_n, b_n and c_n required order by order into the bare action at the begining. Then, such a trick will not be necessary, and we could expect that all divergences will be renormalized into the coupling t and matter fields and other dimensional constants, and that the Wess-Zumino action would not change form.

The difficulty to prove the renormalizability of our model originates from the fact that to preserve diffeomorphism invariance, the Wess-Zumino action should not change form, while the symmetry to preserve the form of the Wess-Zumino action would be nothing but diffeomorphism invariance itself. What can be done seems to be merely to check for any self-consistencies of the theory. In this paper we have shown that our model can preserve diffeomorphism invariance self-consistently up to order t_r^2 (or order t_r^4 for $Z_{\hat{h}}$). Since an extension to higher order would be unique due to the background-metric independence for the traceless mode, it is expected that our model preserves diffeomorphism invariance to all orders.

Here we comment on studies by Kawai, Kitazawa and Ninomiya [30]. While we apply dimensional regularization to the theory $\mathcal{I} = S + I$ (1.3), defined on the metric $\bar{g}$, it seems that, as far as using dimensional regularization, one could apply it to the theory (1.1), defined on the metric g, taking great care with the conformal mode dependence, as discussed by them. In their method it is expected that the coefficients a_r, b_r and c_r are automatically determined, while our method has the advantage that conformal mode dependence is manifest, and hence there is an analogy to 2D quantum gravity. The relation between the two methods is unclear at present. However, it seems that the two methods might be the same in the limit $\epsilon \to 0$, at least up to order t_r^2, because at this order the divergences of the self-energy diagrams of the conformal mode cancel out for arbitrary values of a_r, b_r and c_r. Detailed analyses of this matter seem to be important to prove renormalizability to all orders.

To determine the values a_1, b_1 and c_1, it seems that we must carry out full two-loop calculations. On the other hand, there is a naive argument that the regulator of the measure is given by the bilinear term of quantum fields in the combined Lagranigan, $\mathcal{L} = \frac{1}{2}\Phi^t \mathcal{K} \Phi + o(\Phi^3)$, where $\Phi^t = (\phi, h^\mu{}_\nu, A_\mu)$. Then, using the special property of fourth order operators, we obtain an annomalous contribution formally expressed in the form [16]

$$-2Tr\Big(\delta\omega\, e^{-\varepsilon\mathcal{K}}\Big)\ , \tag{3.80}$$

where ε is a cut-off. This is essentially a one-loop contribution, and hence it determines the values of w_f and w_g. The proof in Ref.[16] of the Wess-Zumino condition for fourth order operators now implies that w_l $(= l'_1) = 0$, which is consistent with what we mentioned above. Here, it is worth commenting on our previous work. In the proof given in section 3 in Ref.[16], it was assumed that an action S satisfying the integrability condition [32] exists. Therefore, the proof implies that, if it does exist, the variation of the part of the Wess-Zumino action S related to one-loop diagrams is given by (3.80). Applying the proof to our model, existence at $t = 0$ is guaranteed by condition (3.14), but not for $t \neq 0$. However, since the $\hat{g}$ dependence of S is determined by the background-metric independence of the conformal mode, the h dependence of S is uniquely determined by the background-metric independence of the traceless mode, as mentioned in the Introduction. Thus, the existence of S is guaranteed in our model, so that $w_l = 0$.

Here we conjecture that only the one-loop diagrams contribute to $\delta_\omega W_{reg}$, and thus $a_1 = 2w_f$ and $b_1 = 2w_g$, because (3.80) is the most simple expression giving the Wess-Zumino action at all orders. Using the generalized Schwinger-DeWitt technique [12], as in Ref.[16], we obtain [11]

$$
\begin{aligned}
2w_f &= -\frac{18a_0^2 + 124a_0b_0 + 109b_0^2}{72b_0} , \qquad (3.81) \\
2w_g &= \frac{14}{9}a_0 + \frac{115}{72}b_0 . \qquad (3.82)
\end{aligned}
$$

Here, u_l is not calculated.

3.4 The t_r^2-order corrections of the cosmological constant

Let us compute the quantum corrections of the cosmological constant, $\Lambda e^{\alpha\phi}$. In 4 dimensions, the constant α should satisfy the equation [13]

$$
\alpha = \dim[\Lambda] = 4 + \gamma_\Lambda , \qquad (3.83)
$$

where γ_Λ is the anomalous dimension. Here, we calculate γ_Λ up to order t_r^2. The contribution to zeroth order of t_r is given by the one-loop diagram in Fig. 1, as in the two-dimensional model. The diagrams of order t_r^2 contributing to the renormalization factor proportional to α^2 are the two 2-loop diagrams, (a) and (b), and the 1-loop diagram, (c), in Fig. 6. Unlike the two-dimensional model, there are contributions proportional to α^3 and α^4. They are given by diagrams (a) and (b) in Fig. 7, respectively.

---

[11] In Ref.[16] we used the gauge-fixing condition that the conformal mode and the traceless mode are mixed. However, this is incorrect, because we did not introduce the coupling t for the conformal mode. To ensure gauge independence, we here use the Feynman-type gauge $\zeta = 1$ in (3.41), and the coefficients a and b are re-calculated. Now, the off-diagonal term in the operator $\mathcal{K}$ becomes fourth order, but it can be managed in the perturbation by reducing the expression of $Tr \log \mathcal{K}$ to the formula given in Ref.[12] by using an identity like, for example $\Box\hat{\nabla}_\mu V_\nu = \hat{\nabla}_\mu \Box V_\nu + 2\hat{R}^\alpha{}_{\nu\mu\beta}\hat{\nabla}^\beta V_\alpha + \frac{1}{4}\hat{R}\hat{\nabla}_\mu V_\nu$, repeatedly, where the constraints $\hat{R}_{\mu\nu} = \frac{1}{4}\hat{g}_{\mu\nu}\hat{R}$ and $\hat{\nabla}_\mu \hat{R} = 0$ are imposed as in Ref.[16], so that the coefficient c_r cannot be determined.


(a) (b) (c)

Fig. 6: Contributions of order t_r^2 proportional to α^2.


(a) (b)

Fig. 7: Contributions of order t_r^2 proportional to α^3 and α^4.

All of the contributions from Fig. 1, the sum of (a) and (b) in Fig. 6, and the other three ones, (c) in Fig. 6 and (a) and (b) in Fig. 7, are proportional to the tadpole-type integral $\int d^D k/k^4$. They thus yield only simple poles, which are summarized as

$$Z_\Lambda - 1 = -\Big(\frac{1}{8\tilde{b}_r} + \frac{7}{32}\frac{\tilde{t}_r^2}{\tilde{b}_r} + \frac{\tilde{t}_r^2}{96}\Big)\frac{\alpha^2}{\epsilon} + \frac{1}{1152}\frac{\tilde{t}_r^2}{\tilde{b}_r}\frac{\alpha^3}{\epsilon} - \frac{1}{3072}\frac{\tilde{t}_r^2}{\tilde{b}_r^2}\frac{\alpha^4}{\epsilon} \ . \tag{3.84}$$

The first term in the parentheses of the r.h.s. comes from the one-loop diagram in Fig. 1, and the second term comes from the sum of the two-loop diagrams, (a) and (b) in Fig. 6. The third in the parentheses is the contribution from the one-loop (but order t_r^2) diagram (c) in Fig. 6. The last two terms on the r.h.s. come from (a) and (b) in Fig. 7, respectively.

There is no counterterm insertion diagrams, because all subdiagrams (namely, the sum of (a) and (b) in Fig. 3, the sum of (a) and (b) in Fig. 8, and (c) itself in Fig. 8) give finite contributions. Furthermore, diagrams (a) and (b) in Fig. 9 each gives a finite contribution. The finiteness of the sum of diagrams (a) and (b) in Fig. 8 and of (c) in Fig. 8 is also evidence of renormalizability.


Fig. 8: Diagrams which give finite contributions I.


Fig. 9: Diagrams which give finite contributions II.

The sum of the diagram of type (a) and associated counterterm-insertion diagrams (b) and (c) in Fig. 10 does not yield simple poles, where the diagram with the gray circle shown in Fig. 11 stands for the diagrams in Figs. 1, 6, 7, 8 and 9. If the gray part is a finite diagram, there is no counterterm of type (c) in Fig. 10, and the sum of the diagrams becomes finite. If the gray part is a divergent diagram, the sum of the diagrams yields only double poles, which only play the role to make an anomalous dimension finite.


Fig. 10: Diagrams that give double poles or finite contributions.


Fig. 11: Symbolical description of Fig. 1 and Figs. 6–9.

Hence, we obtain the anomalous dimension up to order t_r^2 as

$$\gamma_\Lambda = \frac{1}{4}\frac{\alpha^2}{b_r} + t_r^2\Big(\frac{7}{8}\frac{\alpha^2}{b_r} + \frac{\alpha^2}{48} - \frac{1}{288}\frac{\alpha^3}{b_r} + \frac{1}{512}\frac{\alpha^4}{b_r^2}\Big) \ . \tag{3.85}$$

The contributions proportional to α^n with $n > 4$ appear at order t_r^4 and higher.

Until now, we have not specified the value of b_r, which is required by diffeomorphism invariance. Doing so, we obtain

$$\gamma_\Lambda = \frac{1}{4}\frac{\alpha^2}{b_0} + t_r^2\Big(-\frac{b_1}{4}\frac{\alpha^2}{b_0^2} + \frac{7}{16}\frac{\alpha^2}{b_0} + \frac{\alpha^2}{48} - \frac{1}{288}\frac{\alpha^3}{b_0} + \frac{1}{512}\frac{\alpha^4}{b_0^2}\Big) \ . \tag{3.86}$$

Thus, (3.83) becomes biquadratic at order t_r^2.

The cosmological constant should be physically real. Supposing that there is a real solution for the biquadratic equation, it can be solved perturbatively as $\alpha = \sum_n \alpha_n t_r^{2n}$. We then obtain

$$\begin{aligned} \alpha_0 &= 2b_0\Big(1 - \sqrt{1 - \frac{4}{b_0}}\Big) \ , \\ \alpha_1 &= \frac{1}{\sqrt{1-\frac{4}{b_0}}}\Big\{\Big(-\frac{1}{4}\frac{b_1}{b_0^2} + \frac{7}{16}\frac{1}{b_0} + \frac{1}{48}\Big)\alpha_0^2 - \frac{1}{288}\frac{\alpha_0^3}{b_0} + \frac{1}{512}\frac{\alpha_0^4}{b_0^2}\Big\} \ , \end{aligned} \tag{3.87}$$

where we choose the solution of α_0 so that $\alpha_0 \to 4$ in the classical limit, $b_0 \to \infty$.

Let us compare our model with the dynamical triangulation approach to 4D quantum gravity. In this approach there is no ambiguity choosing the measure, unlike with Regge's calculus. Furthermore, although this lattice model has parameters corresponding to the gravitational constant and the cosmological constant, there is no dimensionless coupling t_r, unlike in the usual lattice gauge theories. This indicates that, in continuum theory, the measure of a gravitational field including fourth order parts may be uniquely determined. Here, we estimate the strength of the coupling constant t_r realized in recent "numerical experiments" [21], which suggest that α is imaginary for $N_A = 0$ but becomes real for $N_A \geq 1$.

Recall that b_0 is slightly greater than 4 for a small value of N_A. Thus, α_0 is close to 8. Therefore, the α^3- and α^4-terms and also the third term proportional to α^2 in the parentheses of (3.86) are estimated as having a value on the order $o(1) \times t_r^2$, while the second term in the parentheses, which comes from the sum of the two-loop diagrams (a) and (b) in Fig. 6, is more than 10-times greater than them. Therefore, we neglect the terms proportional to α^3 and α^4. Then, the equation reduces to a quadratic form, and we obtain the following discriminant:

$$J = b_0 + \Big(b_1 - \frac{7}{4}b_0\Big)t_r^2 - 4 \ . \tag{3.88}$$

The results of the simulations [21] can now be described as $J < 0$ for $N_A = 0$, but $J > 4$ for $N_A \geq 1$. If $b_1 = 2w_g$ is assumed,[12] there is a solution of the inequality, and we obtain the relation $0.025 < t_r^2 < 0.040$ for the coupling constant. If b_1 is neglected, the central value shifts by about +0.01. Expression (3.87) is valid in this region.

4 Conclusions and Discussion

In this paper we considered the problems of renormalizability and diffeomorphism invariance in a 4D quantum theory of gravity. We considered a perturbation theory for the traceless mode, where the dimensionless coupling constant t was introduced, while the conformal mode was managed exactly. We found evidence that there is a model that satisfies the two conditions of renormalizability and diffeomorphism invariance simultaneously; it is uniquely determined to be model (3.9) with (3.14) if we do not introduce fourth order fields, except for metric fields. In particular, the vanishing of l_1 and l_1' in (3.25), which is required by the integrability condition discussed by Riegert, Fradkin and Tseytlin [32], gives a breakthrough to the two-loop renormalizability of 4D quantum gravity.

Our conclusion seems to be plausible from the viewpoint of 4D simplicial quantum geometry, because dynamical triangulation chooses a measure uniquely, and therefore the measure of the corresponding continuum theory may be determined uniquely. We expect that it may be given by the fourth order parts of our model.

The fourth order model in 4 dimensions is quite similar to the second order model in two dimensions. We can rather easily compute the quantum corrections of the cosmological constant up to order t_r^2, as in the case of 2D quantum gravity, and we obtain the biquadratic equation (3.83) with (3.86). Quantum corrections of the Einstein-Hilbert term are also quite important. Determining these is a future project.

In this paper we did not consider gauge interactions, such as QED and Yang-Mills theories. If gauge couplings are introduced, one has to add a new coupling, $\kappa\phi Tr(F_{\mu\nu}F^{\mu\nu})$, to the action S. The coefficient κ then becomes a function of the gauge coupling, determined from the non-local logarithmic terms of the regularized vacuum polarizations of gauge fields.

To this point, we have focused only on the two fundamental conditions of renormalizability and diffeomorphism invariance. Theories satisfying these two conditions could be called "quantum geometry" rather than "quantum gravity". To discuss quantum gravity, which should describe our universe, we need to define the S-matrix.

---

[12] Note that $2w_g$ is negative and that $|2w_g|$ is less than b_0 for small N_A.



Fig. 12: Image of the S-matrix of a fire ball/black hole formation and evaporation. The lines with arrows denote matter fields. The wavy lines are gravitational fields in the low-energy asymptotic region. In the fire-ball region, the fourth order terms are turned on.

Recall, as computed in section 3.4, the cosmological constant receives nontrivial quantum corrections, because the propagators of the gravitational modes are proportional to $1/k^4$, and tadpole-type diagrams have logarithmic divergences. This is one difference between the fourth order theories and second order theories merely with negative-metric states. One can define diffeomorphism-invariant correlation functions in such a fully fluctuating geometry. However, when one attempts to define the S-matrix, one encounters the problem of how to define the asymptotic states for the gravitational modes. Since the states are always dressed by the conformal mode when the fourth order kinetic term of the conformal mode exists, it seems that one has to consider an unusual interaction picture in which the fourth order parts are turned off in the asymptotic region, as in a picture of a quantum black hole, where an asymptotically flat region exists (Fig. 12) [38]. In this picture, negative-metric states in the fourth order terms required by diffeomorphism invariance do not appear in the low-energy asymptotic region, because there, gravity is ruled by the Einstein-Hilbert term. Hence, the negative-metric state is unstable and seems to be confined in the internal high-energy region, where the fourth order terms are turned on [39, 9, 40].

More directly, as a consequence of the background-metric independence, or diffeomorphism invariance, the presence of the $\phi\bar{F}$ term in the action implies that we can remove the Weyl term, $\bar{F}$, by shifting the conformal mode by a constant. Thus, although in this "gauge" the theory becomes unrenormalizable, we can remove the negative-metric state for the traceless mode. This situation is somewhat reminiscent the argument of the R_ξ gauge in spontaneously broken gauge theory [41].

Let us compute the degrees of freedom in this case. We must now treat the $\bar{R}^2$ term and the Einstein-Hilbert term. Then, the gauge-fixing term $\chi^\mu\Box\chi_\mu$ in (3.41) is not necessary. Hence, $N_{\mu\nu}$ is proportional to $-\partial_\mu\partial_\nu + m'^2\delta_{\mu\nu}$, where m'^2 originates from the Einstein-Hilbert term, which is proportional to m^2. The ghost determinant is now given by $\det^{1/2}(-\partial_\mu\partial_\nu + m'^2\delta_{\mu\nu}) \det M^{GH}_{\mu\nu}$, where $M^{GH}_{\mu\nu}$

is the usual second order ghost operator of diffeomorphism invariance defined by the relation $M^{GH}_{\mu\nu}\xi^{\nu} = \delta\chi_{\mu}|_{h=0}$. Note that $\det(-\partial_{\mu}\partial_{\nu} + m'^2\delta_{\mu\nu}) = \det(-\Box + m'^2)|_{\text{a scalar}}$ (see, for instance, Ref.[12]), so that the number of ghost degrees of freedom is $4+4+1=9$. Hence, the total number of degrees of freedom becomes $2\times 1+9-9=2$. Thus, the negative-metric state related to the conformal mode is also removed by ghosts. We expect that in the general case, the negative-metric states for the traceless mode may be confined, due to diffeomorphism invariance.

Acknowledgements

I would like to thanks H. Kawai, Y. Kazama and T. Yoneya for helpful comments and suggestions on the manuscript. I also acknowledge H. Hagura, S. Horata, Y. Kitazawa, F. Sugino, A. Tsuchiya and T. Yoshikawa for valuable discussions.

References

[1] B. DeWitt, in Relativity, Groups and Topology, eds. B. DeWitt and C. DeWitt (Gordon and Breach, New York, 1964); Phys. Rev. **160** (1967) 1113; Phys. Rev. **162** (1967) 1195, 1239.

[2] G. 't Hooft and M. Veltmann, Ann. Inst. Henri Poincare **XX** (1974) 69;
M. Veltmann, in Methods in Field Theory, Les Houches 1975.

[3] S. Deser and P. van Nieuwenhuizen, Phys. Rev. Lett. **32** (1974) 69; Phys. Rev. **D10** (1974) 401, 411.

[4] S. Weinberg, in General Relativity, an Einstein Centenary Survay, eds. S. Hawking and W. Israel (Cambridge Univ. Press, Cambridge, 1979);
R. Gastmans, R. Kallosh and C. Truffin, Nucl. Phys. **B133** (1978) 417;
S. Christensen and M. Duff, Phys. Lett. **79B** (1978) 213.

[5] R. Utiyama and B. DeWitt, J. Math. Phys. **3** (1962) 608.

[6] S. Deser, Proceedings of the Coference on Gauge Theories and Modern Field Theories, edited by R. Arnowitt and P. Nath (MIT Press, Cambridge, 1975);
S. Weinberg, Proceedings of the XVIIth International Conference on High Energy Physics, edited by J. R. Smith (Rutherford Laboratory, Chilton, Didcot, 1974).

[7] K. Stelle, Phys. Rev. **D16** (1977) 953; Gen. Rel. Grav. **9** (1978) 353;
A. Salam and J. Strathdee, Phys. Rev. **D18** (1978) 4480.

[8] J. Julve and M. Tonin, Nuovo Cim. **46B** (1978) 137.

[9] E. Tomboulis, Phys. Lett. **97B** (1980) 77;
B. Hasslacher and E. Mottola, Phys. Lett. **99B** (1981) 221.

[10] E. Fradkin and A. Tseytlin, Nucl. Phys. **B201** (1982) 469; Phys. Lett. **104B** (1981) 377.

[11] N. Barth and S. Christensen, Phys. Rev. **D28** (1983) 1876.

[12] A. Barvinsky and G. Vilkovsky, *The generalized Schwinger-DeWitt technique in gauge theories and quantum gravity*, Phys. Rep. **119** (1985) 1.

[13] I. Antoniadis and E. Mottola, Phys. Rev. **D45** (1992) 2013;
S. Odintsov, Z. Phys. **C54** (1992) 531.

[14] I. Antoniadis, P. Mazur and E. Mottola, Nucl. Phys. **B388** (1992) 627.

[15] I. Antoniadis, P. Mazur and E. Mottola, Phys. Lett. **B323** (1994) 284, **B394** (1997) 49;
I. Antoniadis and S. Odintsov, Phys. Lett. **B343** (1995) 76.

[16] K. Hamada and F. Sugino, Nucl. Phys. **B553** (1999) 283.

[17] D. Weingarten, Phys. Lett. **B90** (1980) 285, Nucl. Phys. **B210** [FS6] (1982) 229;
V. Kazakov, Phys. Lett. **150B** (1985) 282;
F. David, Nucl. Phys. **B257** (1985) 45;
J. Ambjørn, B. Durhuus and J. Fröhlich, Nucl. Phys. **B257** (1985) 433;
D. Boulatov, V. Kazakov, I. Kostov and A. Migdal, Nucl. Phys. **B275** (1986) 641.

[18] E. Brezin and V. Kazakov, Phys. Lett. **B236** (1990) 144;
M. Douglas and S. Shenker, Nucl. Phys. **B335** (1990) 635;
D. Gross and A. Migdal, Phys. Rev. Lett. **64** (1990) 127.

[19] J. Ambjørn, B. Durhuus and T. Jonsson, *Quantum Geometry*, (Cambridge Univ. Press, Cambridge, 1997).

[20] M. Agishtein and A. Migdal, Nucl. Phys. **B385** (1992) 395;
J. Ambjørn and J. Jurkiewicz, Phys. Lett. **B278** (1992) 42;
S. Catterall, J. Kogut and R. Renken, Phys. Lett. **B328** (1994) 277.

[21] S. Bilke, Z. Burda, A. Krzywicki, B. Petersson, J. Tabaczek and G. Thorleifsson, Phys. Lett. **B418** (1998) 266; Phys. Lett. **B432** (1998) 279;
H. Egawa, A. Fujitu, S. Horata, N. Tsuda and T. Yukawa, Nucl. Phys. **B (Proc. Suppl.) 73** (1999) 795; *Phase Transition of 4D Simplicial Quantum Gravity with* $U(1)$ *Gauge Field*, Proc. Lattice 99, Pisa (June-July 1999)/hep-lat/9908048.

[22] M. Green, J. Schwarz and E. Witten, *Superstring Theory* (Cambridge Univ. Press, Cambridge, 1978);
J. Polchinski, *String Theory* (Cambridge Univ. Press, Cambridge, 1998).

[23] T. Yoneya, Prog. Theor. Phys. **51** (1974) 1907.

[24] V. Knizhnik, A. Polyakov and A. Zamolodchikov, Mod. Phys. Lett. **A3** (1988) 819.

[25] J. Distler and H. Kawai, Nucl. Phys. **B321** (1989) 509;
F. David, Mod. Phys. Lett. **A3** (1988) 1651.

[26] K. Hamada, Phys. Lett. **B300** (1993) 322;
K. Hamada and A. Tsuchiya, Int. J. Mod. Phys. **A8** (1993) 4897.

[27] J. Russo and A. Tseytlin, Nucl. Phys. **B382** (1992) 259;
A. Chamseddine, Phys. Lett. **B256** (1991) 379; Phys. Lett. **B258** (1991) 97;
S. Odintsov and I. Shapiro, Phys. Lett. **B263** (1991) 183; Int. J. Mod. Phys. **D1** (1993) 571;
S. deAlwis, Phys. Lett. B289 (1992) 278;
A. Bilal and C. Callan, Nucl. Phys. **B394** (1993) 73.

[28] S. Hirano, Y. Kazama and Y. Satoh, Phys. Rev. **D48** (1993) 1687;
Y. Kazama and Y. Satoh, Phys. Rev. **D50** (1994) R2368;
D50 (1994) 3889;
Y. Kazama, Y. Satoh and A. Tsuchiya, Phys. Rev. **D51** (1995) 4265.

[29] H. Kawai and R. Nakayama, Phys. Lett. **B306** (1993) 224;
S. Ichinose, Nucl. Phys. **B457** (1995) 688.

[30] H. Kawai and M. Ninomiya, Nucl. Phys. **B336** (1990) 115;
H. Kawai, Y. Kitazawa and M. Ninomiya, Nucl. Phys. **B393** (1993) 280; Nucl. Phys. **B404** (1993) 684;
T. Aida, Y. Kitazawa, H. Kawai and M. Ninomiya, Nucl. Phys. **B427** (1994) 158;
T. Aida, Y. Kitazawa, J. Nishimura and A. Tsuchiya, Nucl. Phys. **B444** (1995) 353;
H. Kawai, Y. Kitazawa and M. Ninomiya, Nucl. Phys. **B467** (1996) 313;
T. Aida and Y. Kitazawa, Nucl. Phys. **B491** (1997) 427.

[31] J. Wess and B. Zumino, Phys. Lett. **37B** (1971) 95;
S. Treiman, R. Jackiw, B. Zumino and E. Witten, *Current Algebra and Anomalies* (World Scientific Publishing 1985).

[32] R. Riegert, Phys. Lett. **134B** (1984) 56;
E. Fradkin and A. Tseytlin, Phys. Lett. **134B** (1984) 187.

[33] M. Duff, Nucl. Phys. **B125** (1977) 334.

[34] A. Polyakov, Phys. Lett. **103B** (1981) 207;
K. Fujikawa, Phys. Rev. **D25** (1982) 2584;
O. Alvarez, Nucl. Phys. **B216** (1983) 125;
E. D'Hoker and D. Phong, Rev. Mod. Phys. **60** (1988) 917.

[35] L. Abbott, Nucl. Phys. **B185** (1981) 189.

[36] S. Deser, M. Duff and C. Isham, Nucl. Phys. **B111** (1976) 45.

[37] C. Itzykson and J. Zuber, *Quantum Field Theory*, (McGraw-Hill Book Co, 1980).

[38] S. Hawking, Comm. Math. Phys. **43** (1975) 199; Phys. Rev. **D14** (1976) 2460.

[39] T. Lee and G. Wick, Nucl. Phys. **9** (1969) 209.

[40] M. Kaku, Nucl. Phys. **B203** (1982) 285;
D. Boulware, G. Horowitz and A. Strominger, Phys. Rev. Lett. **50** (1983) 1726;
E. Tomboulis, Phys. Rev. Lett. **52** (1984) 1173; Phys. Lett. **B389** (1996) 225.

[41] K. Fujikawa, B. Lee and A. Sanda, Phys. Rev. **D6** (1972) 2923.